


Iowa Toxics Sampling 2004

Results For Selected Pollutants



Air Quality Bureau
Iowa Department of Natural Resources

Table of Contents

[Summary \(7 pages\)](#)

[Toxic Monitoring Network Description](#)

[Toxic Monitoring Sites Map](#)

[Cedar Rapids Site Aerial Photo](#)

[Davenport Site Aerial Photo](#)

[Des Moines Site Aerial Photo](#)

[Cancer Risk Summary](#)

[Concentration Summary](#)

[Percent Data Capture](#)

[Annual Precision Statistics](#)

[Cancer Risks over EPA Benchmark Chart](#)

[Time Response of Formaldehyde Concentrations](#)

[Time Response of Acetaldehyde Concentrations](#)

[Time Response of Benzene Concentrations](#)

[Raw Data - Formaldehyde](#)

[Raw Data - Acetaldehyde](#)

[Raw Data - Benzene](#)

Summary

During the 2004 calendar year the Department operated three toxic monitoring sites in three Iowa cities, selected either because they were heavily populated or heavily industrialized. At each site, two types of toxic samples were taken. Air samples were taken in specially treated stainless steel canisters, in order to capture volatile organic compounds (VOC's) with low molecular weight. Air was also sampled through cartridges in order to investigate the concentration of carbonyl compounds. Both of types of samples were prepared and analyzed at the University of Iowa air toxics analysis lab in Iowa City. The canisters were analyzed according to two different protocols. The EPA's speciated non-methane organic compound (SNMOC) protocol was used to analyze the canister contents using a gas chromatograph and flame ionization detector in order to identify compounds known to be involved in the formation of ozone. EPA's TO-15 protocol was also used to analyze the canister contents with a gas chromatograph and mass spectrometer (GCMS) to quantify low molecular weight toxics commonly found in urban air and identified on EPA's Urban Air Toxics Monitoring Program, (UATMP) list of target compounds. Analysis of carbonyl cartridge extracts was performed using High Performance Liquid Chromatography and an ultraviolet detector according to EPA's method TO-11A. This report includes data from three toxic pollutants (formaldehyde, benzene, and acetaldehyde) that have historically exceeded levels the EPA has determined would pose a greater than one in one million increased cancer risk for a lifetime of exposure.

Sample Collection

Monitors were operated in Des Moines, Cedar Rapids, and Davenport. Toxic samples were collected by the Local County Health Departments in Des Moines and Cedar Rapids; Davenport samples were collected by the University of Iowa Hygienic Laboratory.

Scope

Section 112 of the Clean Air Act identifies 188 hazardous air pollutants (HAPS) produced in sufficient quantities to warrant regulatory action. EPA has selected 32 of these HAP's along with diesel particulate matter for a periodic nation-wide review, the national air toxics assessment (NATA). Of the 33 pollutants assessed in the NATA, 17 were quantified in this study, 15 on the TO-15 list and 2 on the TO-11A list. Toxic metals (quantified using EPA method IO 3.5) and Polycyclic Aromatic Hydrocarbons (quantified using EPA method TO-13A) were not measured by this study, nor were other toxics such as mercury and dioxin. Data in this report is limited to two carbonyl compounds, formaldehyde and acetaldehyde and one volatile organic compound, benzene. These three compounds have historically been present in Iowa at levels greater than the EPA determined one in one million increase in cancer risk for a lifetime exposure.

Sampling Schedules

Samples were gathered on a schedule of one sample every twelfth day. Every sixth day monitoring for carbonyl compounds was conducted at the Des Moines site during the ozone season (April through October), and at Cedar Rapids and Davenport sites from July to September. If a scheduled sample was missed, an unscheduled sample was substituted for the missing data point if that sample was taken before the next scheduled sampling day. In calculations of average pollutant levels and cancer risk the additional samples that were taken during summer time were averaged to estimate the one in twelve average and avoid introduction of a seasonal bias to the data.

Data Capture

The data capture rate is defined as the ratio of the number of samples taken (including scheduled and valid substitute samples) divided by the number of scheduled samples.

Data Handling

All data used in this report was measured at concentrations above the method detection limit (MDL). It was not necessary to perform any data substitutions when measured concentrations below the MDL.

This report characterizes only the cancer risk associated with exposure to the toxic contaminants measured, and does not quantify other “non-cancer” risks such as neurological or reproductive damage associated with the measured exposure levels. The cancer risk associated with a given exposure level was quantified only when an Air Unit Cancer Risk was available in EPA’s Integrated Risk Information System (IRIS) database. Pollutants were selected for inclusion in this report, based on the screening criteria that the excess cancer risk resulting from a lifetime exposure to the average contaminant concentration measured was greater than the EPA benchmark of one in a million excess risk.

Precision Data

Precision data are reported for the total number of collocated pairs of canisters or cartridges collected. Precision statistics shown in this report have been calculated according to 40 CFR Part 58, Appendix A using the methodology applicable to collocated fine particulate data pairs.

Results of the Analysis

Formaldehyde, acetaldehyde, and benzene were measured at levels above the EPA benchmark at all Iowa sites. Formaldehyde levels measured during the study period are associated with a much higher cancer risk than any other pollutant measured in this study.

IRIS specifies different levels of certainty associated with its cancer risk factors. Benzene is classified as a known human carcinogen (Class A). Formaldehyde is a Class B1 carcinogen, and acetaldehyde is classified as a Class B2 carcinogen. Class B contains probable human carcinogens; Class B1 pollutants are associated with limited evidence of carcinogenicity in humans but sufficient evidence of carcinogenicity in animals, whereas a B2 classification indicates only sufficient evidence of carcinogenicity in animals (1).

A primary contaminant is directly emitted into the ambient air from its source. A secondary contaminant is formed from a chemical reaction of other contaminants already present in the atmosphere from natural or anthropogenic sources.

Benzene is a primary contaminant, with emissions largely attributed to vehicular traffic. Formaldehyde and acetaldehyde are both primary and secondary contaminants. Motor vehicle emissions contribute to primary emissions by incomplete combustion of fuel; secondary formation results from photochemical oxidation of exhaust pipe pollutants. Secondary formation of these pollutants is enhanced in the summertime due to suitable weather conditions such as higher temperature and greater hours of sunlight. Formaldehyde is also produced in large quantities by natural events such as forest or brush fires (2).

In interpreting the results of risk assessment contained in this type of report, EPA has encouraged States to compare the risks caused by toxic outdoor air pollution to other risks experienced in everyday life. The highest excess lifetime cancer risk identified in this report is 3.1 excess cancers per 100,000 people (3.1×10^{-5}), associated with average measured formaldehyde levels in the outdoor air at the urban Davenport monitoring site. For comparison, the lifetime risk of dying in a car accident is a 1.7×10^{-2} , or almost 600 times higher, and the lifetime risk of being killed by lightning is 1.2×10^{-5} , or approximately 2 -1/2 times less than developing cancer at this level of formaldehyde exposure (3).


References:

1. Integrated Risk Information System: <http://www.epa.gov/iris>
2. Canada EPA: <http://www.ec.gc.ca>
3. <http://www.nsc.org/lrs/statinfo/odds.htm>

Air Toxics Monitoring Network 2004


Site ID	Site Label	City	Address	County
191130037	Cedar Rapids, Army Reserve	Cedar Rapids	1599 Wenig Rd. NE	Linn
191530030	Des Moines, Public Health Bldg	Des Moines	19th St. & Carpenter St.	Polk
191630015	Davenport, Jefferson Elementary	Davenport	10th St. & Vine	Scott

Iowa Toxics Monitoring Network 2004



[Back](#)

Cedar Rapids Toxics Monitoring 2004



I-380

Army Reserve

0 1 2 3 Miles



[Back](#)

Davenport Toxics Monitoring 2004

I-280

Jefferson Elementary

0 1 2 3 Miles

[Back](#)



Des Moines Toxics Monitoring 2004

I-35/80

Public Health
Building

I-235

0

2

4

6

Miles

[Back](#)



Cancer Risk Summary (Excess Cancers Per Million People)			
	Cedar Rapids	Des Moines	Devenport
Site / Pollutant	Army Reserve	Public Health Building	Jefferson Elementary
Formaldehyde	27.0	28.2	31.1
Acetaldehyde	4.8	2.2	3.0
Benzene	3.4	4.2	4.9

Concentration Summary (ppb)			
	Cedar Rapids	Des Moines	Davenport
Site / Pollutant	Army Reserve	Public Health Building	Jefferson Elementary
Formaldehyde	1.76 (+/- 0.32)	1.84 (+/- 0.31)	2.03 (+/- 0.34)
Acetaldehyde	1.34 (+/- 0.28)	0.60 (+/- 0.08)	0.83 (+/- 0.11)
Benzene	0.14 (+/- 0.04)	0.17 (+/- 0.05)	0.20 (+/- 0.05)

**Note: Values indicated are the average concentrations in parts per billion measured at each site in 2004
Data from enhanced summer monitoring at the three sites were averaged to prevent seasonal bias.
Values listed in parentheses represent the 95% Confidence Interval for the mean.**


Percent Data Capture			
	Cedar Rapids	Des Moines	Devenport
Site / Pollutant	Army Reserve	Public Health Building	Jefferson Elementary
Formaldehyde	92%	89%	97%
Acetaldehyde	92%	91%	95%
Benzene	87%	93%	100%

[Back](#)


Annual Toxics Precision Statistics				
Statistic/ Pollutant	Number of Pairs	Coefficient of Variation	Lower 95% Confidence Limit	Upper 95% Confidence Limit
Formaldehyde	26	12.3%	10.1%	16.0%
Acetaldehyde	26	14.1%	11.5%	18.3%
Benzene	32	23.7%	19.7%	29.9%

Note: Statistics generated from collocated sample pairs. CV and Confidence Limits calculated according to 40 CFR Part 58 (PM 2.5 method).


Air Toxics With Cancer Risk Over The EPA Benchmark - 2004



Formaldehyde Cancer Risk In Iowa In 2004




Acetaldehyde Cancer Risk In Iowa In 2004



[Back](#)

Benzene Cancer Risk In Iowa In 2004



[Back](#)

Raw Data - Formaldehyde

(Concentration in ppb)

Date	Cedar Rapids	Des Moines	Davenport
	Army Reserve	Public Health Building	Jefferson Elementary
1/10/2004	1.5	0.99	1.3
1/22/2004	0.57	0.62	0.64
2/3/2004	1.1	0.86	
2/4/2004			1.4
2/15/2004	1.5		
2/27/2004	1	1.8	1.3
3/10/2004	1.2	1.6	1.5
3/22/2004	1.4		1.7
4/3/2004	1.4	2.1	1.7
4/9/2004		2.8	
4/15/2004	2.5	4.2	3.2
4/21/2004		2.2	
4/27/2004	1.5	2.7	1.8
5/3/2004		2.6	
5/9/2004	2.5	3	3
5/15/2004		3.1	
5/21/2004		2.2	2
6/2/2004	1.2	1.5	1.4
6/8/2004		2.3	
6/14/2004	2.6	2.3	2.7
6/20/2004		1.5	
6/26/2004	2.2	2	2.8
7/2/2004	3.9	2.5	3.7
7/8/2004	2	1.8	2.5
7/14/2004	2.5	2.3	2.7
7/20/2004	5.2	4.1	4.7
7/26/2004	2.6	2.3	2.9
8/1/2004	3.1	2.1	3.7
8/7/2004		2.4	2.8
8/13/2004	1.8	1.8	2
8/19/2004	1.5	1.6	2
8/25/2004	1.6	1.3	1.5
8/31/2004	1.6	2	2.8
9/6/2004	1.9	0.98	2.2
9/12/2004	3.7	3	3.9
9/18/2004	2.6	2.2	2.9
9/24/2004	1.9	2.1	2.5
9/30/2004	2.4	2.4	2.7
10/6/2004	2.3		2.5
10/12/2004		1.7	1.7
10/18/2004	1.1		
10/21/2004			
10/24/2004			
10/30/2004			1.2
11/5/2004	1.2		1.7
11/17/2004		1	1.4
11/23/2004	0.75		
11/29/2004		0.77	0.93
12/5/2004	0.88		
12/11/2004	0.66	0.72	0.81
12/23/2004	0.88	0.9	1

Raw Data - Acetaldehyde

(Concentration in ppb)

Date	Cedar Rapids Army Reserve	Des Moines Public Health Building	Davenport Jefferson Elementary
1/10/2004	1.5	0.51	0.81
1/22/2004	0.36	0.16	0.34
2/3/2004	0.71	0.27	
2/4/2004			1
2/15/2004	0.77		
2/27/2004	0.89	0.59	0.82
3/10/2004	0.93	0.4	0.63
3/22/2004	0.91		0.61
4/3/2004	0.91	0.53	0.62
4/9/2004		0.84	
4/15/2004	1.8	1	1.1
4/21/2004		0.47	
4/27/2004	1	0.51	0.52
5/3/2004		0.69	
5/9/2004	1.9	0.7	0.88
5/15/2004		0.91	
5/21/2004		0.69	1.3
6/2/2004	0.99	0.65	0.46
6/8/2004		0.5	
6/14/2004	2	0.9	0.85
6/20/2004		0.58	
6/26/2004	1.9	0.74	0.96
7/2/2004	2.4	0.7	1.5
7/8/2004	1.3	0.53	0.85
7/14/2004	1.8	0.83	0.91
7/20/2004	3.8	0.9	1.2
7/26/2004	1.7	0.89	1.1
8/1/2004	2.8	0.69	1.1
8/7/2004		0.67	0.93
8/13/2004	1.4	0.56	0.67
8/19/2004	1.6	0.55	0.54
8/25/2004	1.3	0.42	0.45
8/31/2004	1.3	0.88	1.1
9/6/2004	1.6	0.37	0.63
9/12/2004	3.9	0.77	1.7
9/18/2004	2	0.72	1
9/24/2004	1.8	0.95	1
9/30/2004	2.2	0.73	1.6
10/6/2004	2.2	0.85	1.2
10/12/2004		0.94	1
10/18/2004	0.85		
10/21/2004			
10/24/2004			
10/30/2004			0.47
11/5/2004	0.98	0.57	0.86
11/17/2004		0.37	0.85
11/23/2004	0.5		
11/29/2004		0.27	0.3
12/5/2004	0.57		
12/11/2004	0.51	0.44	0.32
12/23/2004	0.59	0.42	0.46

Raw Data - Benzene

(Concentration in ppb)

Date	Cedar Rapids	Des Moines	Davenport
	Army Reserve	Public Health Building	Jefferson Elementary
1/10/2004	0.31	0.34	0.31
1/22/2004	0.13	0.26	0.13
2/3/2004	0.2	0.36	
2/4/2004			0.36
2/15/2004	0.24		0.26
2/27/2004	0.28	0.27	0.39
3/10/2004	0.19	0.2	0.2
3/22/2004	0.19		0.19
4/3/2004	0.11	0.25	0.13
4/9/2004		0.13	
4/15/2004	0.14		0.19
4/27/2004	0.09	0.18	0.11
5/9/2004	0.09	0.14	0.12
5/21/2004	0.09	0.17	0.27
6/2/2004	0	0	0
6/14/2004	0	0.16	0
6/26/2004	0.06	0.13	0.17
7/8/2004	0.08	0.11	0.14
7/20/2004	0.22	0.38	0.23
8/1/2004	0.15	0.26	0.28
8/13/2004	0.12	0.28	0.25
8/25/2004	0.18	0.38	0.19
9/6/2004	0	0.02	0.08
9/18/2004	0.1	0.2	0.19
9/30/2004	0.27	0.12	0.5
10/12/2004		0.12	0.24
10/21/2004	0.09		
10/24/2004		0.7	0.27
11/5/2004		0	0
11/17/2004		0	0.58
11/29/2004		0	0.16
12/11/2004	0	0	0.09
12/23/2004	0.12	0.12	0.16