

**Iowa Toxics Sampling 2011  
Results for Benzene, Acetaldehyde, and Formaldehyde**



**Air Quality Bureau  
Iowa Department of Natural Resources**

## Table of Contents

Summary: Scope .....	1
Sampling Schedules.....	1
Data Capture .....	1
Data Handling.....	1
Precision Data .....	1
Results of the Analysis .....	1
References .....	2
Air Toxics Monitoring Network 2011 .....	2
Iowa 2011 Air Toxics Monitoring Network .....	3
Cedar Rapids Toxics Monitoring Site .....	4
Davenport Toxics Monitoring Site .....	5
Des Moines Toxics Monitoring Site.....	6
Cancer Risk Summary - Aldehydes (Excess Cancers per Million People).....	6
Excess Cancer Risks per 1 Million People, Aldehydes – 2011 .....	7
Concentration Summary – Aldehydes (ppb).....	7
Cancer Risk Summary - Benzene (Excess Cancers per Million People).....	7
Excess Cancer Risks per 1 Million People, Benzene – 2011 .....	8
Concentration Summary – Benzene (ppb).....	8
Percent Data Capture.....	8
Annual Toxics Precision Statistics .....	9
Formaldehyde Cancer Risk 2011.....	9
Acetaldehyde Cancer Risk 2011 .....	10
Benzene Cancer Risk 2011 .....	11
Raw Data - Formaldehyde.....	11
Raw Data – Acetaldehyde .....	13
Raw Data – Benzene .....	14
Appendix A. Precision Calculations.....	16

## **Summary: Scope**

Section 112 of the Clean Air Act [1] contains the federal strategy for protecting the public from air toxics emissions. The Act specifies a particular list of air toxics called “hazardous air pollutants” (HAPs) for regulatory action [2]. Emitters of large amounts of these HAPs are subject to regulations that require adoption of work practices or installation of control technologies in order to reduce HAP emissions [3]. The Act requires a periodic assessment of the residual health risk posed by the HAPs [4] and adoption of additional control standards where necessary [5].

In order to establish long term trends in HAP concentrations across the nation as a component of its residual risk assessment, the Environmental Protection Agency (EPA) has funded national air toxics trends stations (NATTS) [6]. These sites contain a standard suite of samplers and analytical protocols [7]. Unlike NATTS sites, Iowa’s population-oriented air toxics sites do not have instrumentation to measure toxic metals, polycyclic aromatic hydrocarbons, or black carbon.

A review of the historical air toxics monitoring dataset [8] argues that benzene, formaldehyde, 1,3- butadiene, acrolein, arsenic, hexavalent chromium, and diesel particulate pose the greatest risk to the public health on a national level. Only two of the seven national risk drivers are quantified by the limited air toxics sampling currently conducted in Iowa.

## **Sampling Schedules**

Samples were gathered on a nominal schedule of one sample every twelfth day. In calculations of average pollutant levels and cancer risks, samples collected on a more frequent schedule were averaged over the twelve day period between scheduled samples to estimate a one in twelve sampling schedule and avoid introduction of bias to the data. The monitoring schedule for formaldehyde and acetaldehyde was accelerated to one in six days during ozone season (April through October).

## **Data Capture**

The data capture rate is defined as the ratio of the number of samples taken (including scheduled and valid substitute samples) divided by the number of scheduled samples in the year. Due to the relocation of the toxics analysis laboratory, the data capture rate for benzene for all three sites was 50% during the first quarter of 2011. EPA data analysis guidelines usually require 75% data completeness across each sampling quarter.

## **Data Handling**

This report characterizes only the cancer risk associated with exposure to the toxic contaminants measured, and does not quantify other “non-cancer” risks such as neurological or reproductive damage associated with the measured exposure levels. The cancer risk associated with a given exposure level was quantified only when an Air Unit Cancer Risk was available in EPA’s Integrated Risk Information System (IRIS) database. Pollutants were selected for inclusion in this report, based on the screening criteria that the excess cancer risk resulting from a lifetime exposure to the average contaminant concentration measured was greater than the EPA benchmark of one in a million excess risks. When calculating the cancer risks and annual summary statistics for the selected pollutants, reported data values less than the method detection limit (MDL) are replaced with data values equal to half the MDL. Only Benzene had reported values under the MDL in 2011.

## **Precision Data**

Precision data are reported for the total number of collocated pairs of canisters or cartridges collected. Precision statistics shown in this report have been calculated according to 40 CFR Part 58, Appendix A (2006) using the methodology applicable to collocated fine particulate data pairs. (See Appendix A)

## **Results of the Analysis**

Formaldehyde, acetaldehyde, and benzene concentrations were measured at levels above the EPA benchmark at all Iowa sites. Formaldehyde levels measured during the study period are associated with a much higher cancer risk than any other pollutant measured in this study.

IRIS specifies different levels of certainty associated with its cancer risk factors. Benzene is classified as a known human carcinogen (Class A). Formaldehyde is a Class B1 carcinogen, and acetaldehyde is classified as a Class B2 carcinogen. Class B contains probable human carcinogens; Class B1 pollutants are associated with limited evidence of carcinogenicity

in humans but sufficient evidence of carcinogenicity in animals, whereas a B2 classification indicates only sufficient evidence of carcinogenicity in animals. [9].

A primary contaminant is directly emitted into the ambient air from its source. A secondary contaminant is formed from a chemical reaction of other contaminants already present in the atmosphere from natural or anthropogenic sources.

Benzene is a primary contaminant, with emissions largely attributed to vehicular traffic. Formaldehyde and acetaldehyde are both primary and secondary contaminants. Motor vehicle emissions contribute to primary emissions by incomplete combustion of fuel; secondary formation results from photochemical oxidation of exhaust pipe pollutants. Secondary formation of these pollutants is enhanced in the summertime due to suitable weather conditions such as higher temperature and greater hours of sunlight. Formaldehyde is also produced in large quantities by natural events such as forest or brush fires [10]. In interpreting the results of the risk assessment contained in this type of report, EPA has encouraged States to compare the risks caused by toxic outdoor air pollution to other risks experienced in everyday life. The highest excess lifetime cancer risk identified in this report is 3.3 excess cancers per 100,000 people ( $3.3 \times 10^{-5}$ ), associated with average measured formaldehyde levels in the outdoor air at the urban Des Moines monitoring site. For comparison, the lifetime risk of dying in a car accident is a  $3.3 \times 10^{-3}$ , or approximately 100 times higher, and the lifetime risk of being killed by lightning is  $1.2 \times 10^{-5}$ , or approximately 2.75 times lower than developing cancer at this level of formaldehyde exposure [11].


## References

1. Federal rules regulating air toxics: <http://www.epa.gov/ttn/atw/eparules.html>
2. Current list of HAPs and their health effects: <http://www.epa.gov/ttn/atw/hlthef/hapindex.html>
3. EPA regulations limiting HAPs emissions: <http://www.epa.gov/ttn/atw/mactfnlalph.html>
4. EPA's latest national assessment of health risks due to HAPs: <http://www.epa.gov/ttn/atw/natamain/>
5. Residual risk assessments: <http://www.epa.gov/ttn/atw/rrisk/rtrpg.html>
6. Current list of NATTS sites: <http://www.epa.gov/ttnamti1/files/ambient/airtox/nattsite.pdf>
7. Sampling protocol used to operate NATTS sites: [http://www.epa.gov/ttn/amtic/files/ambient/airtox/NATTS\\_Model\\_QAPP.pdf](http://www.epa.gov/ttn/amtic/files/ambient/airtox/NATTS_Model_QAPP.pdf)
8. Historical review of air toxics monitoring data: <http://www.ladco.org/reports/toxics/sti/>
9. Integrated Risk Information System: <http://www.epa.gov/iris>
10. Reinhardt TE, Ottmar RD. "Baseline Measurements of Smoke Exposure Among Wildland Firefighters." Journal of Occupational and Environmental Hygiene 2004 Sep; 1 (9):593-606. <http://www.ncbi.nlm.nih.gov/pubmed?term=Baseline%20Measurements%20of%20Smoke%20Exposure%20Among%20Wildland>
11. Mortality Odds: [http://www.nsc.org/news\\_resources/injury\\_and\\_death\\_statistics/Documents/Injury\\_Facts\\_37.jpg](http://www.nsc.org/news_resources/injury_and_death_statistics/Documents/Injury_Facts_37.jpg)

## Air Toxics Monitoring Network 2011


Site ID	Site Label	City	Address	County
191130040	Cedar Rapids, Public Health	Cedar Rapids	500 11 <sup>th</sup> St NW	Linn
191530030	Des Moines, Public Health	Des Moines	1907 Carpenter Ave	Polk
191630015	Davenport, Jefferson School	Davenport	10 <sup>th</sup> St & Vine St	Scott

Iowa 2011 Air Toxics Monitoring Network






Cedar Rapids Toxics Monitoring Site






Davenport Toxics Monitoring Site





**Des Moines Toxics Monitoring Site**




**Cancer Risk Summary - Aldehydes (Excess Cancers per Million People)**

<b>Site / Pollutant</b>	<b>Cedar Rapids Public Health</b>	<b>Des Moines Public Health Building</b>	<b>Davenport Jefferson Elementary</b>
Formaldehyde	30.91	33.05	28.68
Acetaldehyde	3.55	2.79	2.82



**Excess Cancer Risks per 1 Million People, Aldehydes – 2011**



**Concentration Summary – Aldehydes (ppb)**

Site / Pollutant	Cedar Rapids Public Health	Des Moines Public Health Building	Davenport Jefferson Elementary
Formaldehyde	2.02 (±0.34)	2.15 (±0.38)	1.87 (±0.37)
Acetaldehyde	0.99 (±0.11)	0.78 (±0.08)	0.78 (±0.10)


Notes: Values indicated are the average concentrations in parts per billion measured at each site in 2011. Data from enhanced summer monitoring at the sites were averaged to prevent seasonal bias. Values listed in parentheses represent the 95% Confidence Interval for the mean.

**Cancer Risk Summary - Benzene (Excess Cancers per Million People)**

Site / Pollutant	Cedar Rapids Public Health	Des Moines Public Health Building	Davenport Jefferson Elementary
Benzene* (with MDL Substitution)	5.37	4.76	5.04
Benzene*	6.05	5.43	5.54

\*IRIS lists two cancer risk estimates for Benzene, and the higher risk estimate is used for the statistics in this report.

**Excess Cancer Risks per 1 Million People, Benzene – 2011**



**Concentration Summary – Benzene (ppb)**

Site / Pollutant	Cedar Rapids Public Health	Des Moines Public Health Building	Davenport Jefferson Elementary
Benzene* (with MDL Substitution)	0.22 (±0.05)	0.20 (±0.03)	0.21 (±0.04)
Benzene*	0.25 (±0.04)	0.22 (±0.02)	0.23 (±0.03)

Notes: Values indicated are the average concentrations in parts per billion measured at each site in 2011. Data from enhanced summer monitoring at the site were averaged to prevent seasonal bias. Values listed in parentheses represent the 95% Confidence Interval for the mean.

**Percent Data Capture**

Site / Pollutant	Cedar Rapids Public Health	Des Moines Public Health Building	Davenport Jefferson Elementary
Formaldehyde	97%	97%	97%
Acetaldehyde	97%	97%	97%
Benzene	87%	84%	71%




### Annual Toxics Precision Statistics


Statistic/ Pollutant	Number of Pairs	Coefficient of Variation	Lower 95% Confidence Limit	Upper 95% Confidence Limit
Formaldehyde	42	2.9%	2.5%	3.6%
Acetaldehyde	42	2.7%	2.3%	3.3%
Benzene	27	15.1%	12.4%	19.6%

Note: Statistics generated from collocated sample pairs. CV and confidence limits calculated according to 2006 methods in Appendix A.

### Formaldehyde Cancer Risk 2011




# Acetaldehyde Cancer Risk 2011





## Benzene Cancer Risk 2011



## Raw Data - Formaldehyde

(Concentration in ppb)

Date	Cedar Rapids Public Health	Des Moines Public Health Building	Davenport Jefferson Elementary
1/3/2011	1.1	1.5	2.1
1/15/2011			
1/27/2011	1.2	1.8	1.3
2/8/2011	0.71	0.84	0.94
2/20/2011	0.87	0.92	1
3/4/2011	1	1.2	1
3/16/2011	2.1	2.1	1.9
3/28/2011	2.1	1.8	1.4
4/3/2011	3.2	2.6	2.2
4/9/2011	2.1	2.2	1.7
4/15/2011	1.4	1.1	1
4/18/2011			1
4/21/2011	1.5	1.4	1.2
4/24/2011	1.5	1.7	1.8

<b>Date</b>	<b>Cedar Rapids Public Health</b>	<b>Des Moines Public Health Building</b>	<b>Davenport Jefferson Elementary</b>
4/27/2011	0.98	1.4	1.1
5/3/2011	1.2	1.9	1.4
5/9/2011	2.7	3.4	
5/15/2011	1.5	1.3	1.3
5/21/2011	2.5	1.9	2.4
5/27/2011	2	2	
5/30/2011			3.6
6/2/2011	2.3	3.7	
6/8/2011	6.5	5.7	6.7
6/14/2011	2.2	1.8	1.8
6/20/2011	2.3		1.9
6/23/2011		1.2	1
6/26/2011	2.3	1.8	1.9
6/29/2011			2.5
7/2/2011	3.5	3	3.8
7/8/2011	3.1	3.8	3.7
7/14/2011	2.8	3.3	
7/20/2011	5.4	5.1	5.6
7/26/2011	3	3.5	3.1
8/1/2011	4.2	6.2	3.2
8/7/2011	1.9	2	1.9
8/10/2011			1.6
8/13/2011	1.7	1.7	
8/19/2011	2.6	2.5	
8/25/2011	1.9	2.8	1.8
8/31/2011	3.3	4.6	
9/6/2011	1.3	2	1.4
9/12/2011	2.8	2.8	3.4
9/15/2011			1.4
9/18/2011	1.8	1.3	1.9
9/21/2011			1.7
9/24/2011	1	1.8	1.4
9/30/2011	1.6	2	1.2
10/6/2011	4.2	4.4	3.5
10/12/2011	1.3	2.2	1.7
10/18/2011	2.3	0.93	0.82
10/24/2011	2.2	2.8	1.2
10/27/2011			1.3
10/30/2011	1.9	2.2	1.6
11/11/2011	1.7	1.7	1.4
11/23/2011	1.2	1.7	1.3
12/5/2011	1.1	1.1	0.72



<b>Date</b>	<b>Cedar Rapids Public Health</b>	<b>Des Moines Public Health Building</b>	<b>Davenport Jefferson Elementary</b>
12/17/2011	1.1	1.2	1.2
12/29/2011	1.9	1.7	1.4

**Raw Data – Acetaldehyde**  
(Concentration in ppb)

<b>Date</b>	<b>Cedar Rapids Public Health</b>	<b>Des Moines Public Health Building</b>	<b>Davenport Jefferson Elementary</b>
1/3/2011	0.74	0.69	1.1
1/15/2011			
1/27/2011	0.64	0.8	0.73
2/8/2011	0.36	0.37	0.43
2/20/2011	1	0.45	0.49
3/4/2011	0.81	0.66	0.56
3/16/2011	1.3	0.87	1
3/28/2011	0.82	0.54	0.42
4/3/2011	1.2	0.87	0.83
4/9/2011	0.9	0.73	0.63
4/15/2011	0.73	0.51	0.47
4/18/2011			0.3
4/21/2011	0.72	0.57	0.44
4/24/2011	1.9	0.63	0.61
4/27/2011	0.44	0.39	0.38
5/3/2011	2.7	0.53	0.46
5/9/2011	1.4	1	
5/15/2011	0.71	0.41	0.36
5/21/2011	0.98	0.61	0.76
5/27/2011	0.74	0.72	
5/30/2011			0.9
6/2/2011	1.4	0.94	
6/8/2011	1.4	1.2	1.3
6/14/2011	0.87	0.67	0.55
6/20/2011	0.95		0.69
6/23/2011		0.38	0.37
6/26/2011	1.2	0.59	0.54
6/29/2011			0.89
7/2/2011	1.1	0.97	0.95
7/8/2011	1.4	1.1	1.3
7/14/2011	1	0.77	
7/20/2011	1.2	0.96	1.2
7/26/2011	0.91	1	1.1
8/1/2011	1.1	0.99	0.85
8/7/2011	1.1	0.59	0.63

Date	Cedar Rapids Public Health	Des Moines Public Health Building	Davenport Jefferson Elementary
8/10/2011			0.52
8/13/2011	0.6	0.51	
8/19/2011	1.2	0.93	
8/25/2011	0.9	0.87	0.64
8/31/2011	1.4	1.2	
9/6/2011	0.59	0.79	0.49
9/12/2011	0.93	0.97	1.4
9/15/2011			0.64
9/18/2011	1	0.54	0.91
9/21/2011			0.76
9/24/2011	1.1	0.77	0.71
9/30/2011	0.7	0.88	0.65
10/6/2011	1.7	1.3	1.7
10/12/2011	1.4	1.2	0.96
10/18/2011	1.3	0.63	0.8
10/24/2011	0.99	1.2	0.53
10/27/2011			1.2
10/30/2011	1	1	1.3
11/11/2011	1.2	1.1	1.3
11/23/2011	0.71	1	0.88
12/5/2011	0.42	0.45	0.54
12/17/2011	0.6	0.81	0.9
12/29/2011	1.4	0.82	0.92

**Raw Data – Benzene**  
(Concentration in ppb)

Date	Cedar Rapids Public Health	Des Moines Public Health Building	Davenport Jefferson Elementary
1/3/2011			
1/15/2011			
1/27/2011			
2/8/2011			
2/20/2011	0.18*	0.22	0.21
3/4/2011	0.23	0.24	0.29
3/16/2011	0.51	0.29	0.14*
3/22/2011	0.22	0.26	0.26
3/28/2011	0.17*	0.19*	0.14*
4/3/2011	0.23	0.19*	0.21
4/9/2011	0.26	0.22	0.28
4/21/2011	0.28	0.25	0.27
4/27/2011	0.15*	0.15*	0.13*

Date	Cedar Rapids Public Health	Des Moines Public Health Building	Davenport Jefferson Elementary
5/3/2011	0.15*	0.17*	0.17*
5/15/2011	0.15*	0.12*	0.09*
5/21/2011			0.19*
5/27/2011	0.17*	0.24	
5/30/2011			0.18*
6/2/2011	0.26	0.25	
6/8/2011	0.21	0.19*	0.19*
6/14/2011		0.16*	0.16*
6/20/2011	0.15*		0.17*
6/23/2011			0.13*
6/26/2011	0.14*	0.15*	0.14*
7/2/2011	0.2*	0.27	0.19*
7/14/2011	0.2*	0.19*	
7/26/2011	0.23	0.31	0.25
8/7/2011	0.17*	0.21	0.16*
8/19/2011	0.31		
8/31/2011	0.25	0.33	
9/12/2011	0.23	0.27	
9/24/2011	0.33	0.29	
9/30/2011		0.17*	0.13*
10/6/2011	0.27	0.22	0.25
10/12/2011			0.34
10/18/2011	0.12*	0.12*	0.16*
10/30/2011	0.18*	0.17*	0.31
11/11/2011	0.45	0.2*	0.32
11/23/2011	0.46	0.31	0.35
12/5/2011	0.19*	0.18*	0.25
12/11/2011			0.42
12/17/2011	0.15*	0.17*	0.24
12/29/2011	0.49	0.3	0.3

\*Indicates sample value reported is less than the minimum detectable limit



## Appendix A. Precision Calculations

Let  $c_i^1$  and  $c_i^2$  represent two concentrations from a particular monitoring location taken on the same day. If both are greater than the MDL, then they may be used to estimate the precision of the data at the sampling location as follows:

First compute the average:

$$\bar{c}_i = \frac{c_i^1 + c_i^2}{2}$$

And the mean difference:

$$d_i = \frac{c_i^1 - c_i^2}{c_i}$$

Define the coefficient of variation for the pair of samples as:

$$CV_i = \frac{d_i}{\sqrt{2}}$$

Compute the root mean square of the individual coefficients of variation to determine the coefficient of variation of the data at the site for the entire year:

$$CV = \sqrt{\frac{\sum_{i=1}^n CV_i^2}{n}}$$

Finally, compute confidence limits in the usual way:

$$\text{Lower Confidence Limit} = CV = \sqrt{\frac{n}{X_{(.05,n)}^{-1}}}$$

$$\text{Upper Confidence Limit} = CV = \sqrt{\frac{n}{X_{(.95,n)}^{-1}}}$$

Where  $X^{-1}$  represents the inverse of the chi-squared distribution.