How You Can Help Protect Our Bees

- Follow integrated pest management practices to first evaluate if a pesticide application is really necessary. If the damage has already been done and is largely cosmetic, treating will not improve the plant's health or appearance.
- Read each pesticide label to determine if it is toxic to bees. The Environmental Hazards Statement will list animals that may be harmed from an application of that product.

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to lakes, streams or ponds. Do not dump rinse water into sewers or other bodies of water. This pesticide is highly toxic to bees. Do not apply this product when bees are active. Apply this product only as specified on this label.

MPORTANT: READ BEFORE USE Read the entire Directions for Use of liability before using this or list and the strength of the product only as specified on the label.

d product container at once. By using conditions, disclaimer of warrantie

 Purchase formulations that are less toxic to bees. Granules and emulsifiable concentrates are safer to use than wettable powders, dusts, or microencapsulated products.

terms are not accer

- 4. Avoid applying any pesticides (especially insecticides and fungicides) to any blooming plant. Bees and other pollinators may be harmed if they contact the pesticide or if they consume nectar or pollen containing pesticides.
- 5. Apply pesticides only after flower petals have fallen if the plant is in need of pest protection.
- If using a white grub insecticide with imidacloprid, mow the lawn within a week of the application to minimize exposure to bees visiting clovers.
- 7. Use products that are less toxic to bees, such as insecticidal soap or *Bacillus thuringiensis*.



- 8. Avoid using any systemic insecticide on plants even after blooming if that product lasts until next season's blooming. Imidacloprid is one of those products that persists, but dinotefuran is active only during the current season.
- When purchasing ornamental plants that are known to attract pollinators, try to purchase ones that do not have insect and plant pathogen pests.
- 10. Learn more online:

www.beeinformed.org
www.entomology.umn.edu/cues/pollinators
www.npic.orst.edu
www.pesticidestewardship.org
www.pollinator.org
www.xerces.org

Prepared by L Jesse (Plant & Insect Diagnostic Clinic), and D Lewis, J Coats, and M Shour (Department of Entomology).


Questions? Contact Iowa State University Extension and Outreach.

...and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, pare proposed in the programs of the programs of the programs. Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue SW, Washington, DC 20250-9410, or call 800-795-3272 (voice) or 202-720-6382 (TDD). USDA is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa.

SP 455 December 2013

IOWA STATE UNIVERSITY Extension and Outreach



Honey bees, bumble bees, orchard bees, mason bees, and various other insects pollinate our fruit and vegetable plants, as well as ornamentals and native plants. Pollinators are critical for our environment and economy, and they need our help!



ollinators, particularly honey bees, are in decline. No one factor is believed to be the cause of bee losses; rather it appears to be a combination of problems (e.g., habitat loss, poor diet, parasites, and pesticide exposure).

By design, insecticides are toxic to insects. Of primary concern are honey bees, bumble bees, native solitary bees, and other beneficial insects, and we can lessen the impacts to them by reducing exposure to these pesticides.


Neonicotinoids are a group of insecticides that have been under greater scrutiny. Imidacloprid is the most widely available neonicotinoid used by homeowners. Other neonicotinoid active ingredients are acetamiprid, clothianidin, dinotefuran, thiacloprid, and thiamethoxam.



Neonicotinoids are being closely examined for several reasons:

- 1. they are a relatively new type of insecticide that has quickly attained worldwide use;
- they are systemic, meaning they move throughout the plant into various tissues, including pollen and nectar, and can be a source of exposure to pollinators;
- 3. they can be present in the ecosystem for a long time; and
- 4. they are highly toxic to bees.

An insecticide does not have to kill in order to have a negative impact. Recent research suggests that exposure to neonicotinoids may affect a bee's ability to fly and navigate, learn, and reproduce.



honey bee

A careful application of imidacloprid to protect your ash tree from emerald ash borer will have minimal risks to pollinators since this tree is windpollinated and the flowers bloom early in the spring and are not attractive to pollinating insects.

Applications to protect a linden tree from Japanese beetles, however, produce a very high risk to bees and other insects. Iowa State University Extension and Outreach does not recommend using imidacloprid just prior to or during bloom since it is so toxic to bees.



